arXiv:1606.05520v2 [math.CA] 18 Aug 2016

Complete monotonicity of some entropies

Ioan Raşa
Department of Mathematics, Technical University of Cluj-Napoca,
Memorandumului Street 28,
400114 Cluj-Napoca,
Romania, ioan.rasa@math.utcluj.ro

Abstract

It is well-known that the Shannon entropies of some parameterized probability distributions are concave functions with respect to the parameter. In this paper we consider a family of such distributions (including the binomial, Poisson, and negative binomial distributions) and investigate the Shannon, Rényi, and Tsallis entropies of them with respect to the complete monotonicity.
keywords: entropies; concavity; complete monotonicity; inequalities
subject class: 94A17; 60E15; 26A51

1 Introduction

Let $c \in \mathbb{R}$, $I_c := \left[0, -\frac{1}{c}\right]$ if $c < 0$, and $I_c := [0, +\infty)$ if $c \geq 0$.

For $\alpha \in \mathbb{R}$ and $k \in \mathbb{N}_0$ the binomial coefficients are defined as usual by

$$\binom{\alpha}{k} := \frac{\alpha(\alpha-1)\dots(\alpha-k+1)}{k!} \quad \text{if } k \in \mathbb{N}, \text{ and } \binom{\alpha}{0} := 1.$$

Let $n > 0$ be a real number such that $n > c$ if $c \geq 0$, or $n = -cl$ with some $l \in \mathbb{N}$ if $c < 0$.

For $k \in \mathbb{N}_0$ and $x \in I_c$ define

$$p_{n,k}^{[c]}(x) := (-1)^k \binom{-\frac{n}{c}}{k} (cx)^k (1+cx)^{-\frac{n}{c}-k}, \quad \text{if } c \neq 0,$$

$$p_{n,k}^{[0]}(x) := \lim_{c\to 0} p_{n,k}^{[c]}(x) = \frac{(nx)^k}{k!}e^{-nx}.$$

Details and historical notes concerning these functions can be found in [3], [7], [21] and the references therein. In particular,

$$\frac{d}{dx}p_{n,k}^{[c]}(x) = n\left(p_{n+c,k-1}^{[c]}(x) - p_{n+c,k}^{[c]}(x)\right). \tag{1}$$

Moreover,

$$\sum_{k=0}^{\infty} p_{n,k}^{[c]}(x) = 1; \tag{2}$$

$$\sum_{k=0}^{\infty} k p_{n,k}^{[c]}(x) = nx, \tag{3}$$

so that $\left(p_{n,k}^{[c]}(x)\right)_{k\geq 0}$ is a parameterized probability distribution. Its associated Shannon entropy is

$$H_{n,c}(x) := -\sum_{k=0}^{\infty} p_{n,k}^{[c]}(x)\log p_{n,k}^{[c]}(x),$$

while the Rényi entropy of order 2 and the Tsallis entropy of order 2 are given, respectively, by (see [18], [20])

$$R_{n,c}(x) := -\log S_{n,c}(x); \quad T_{n,c}(x) := 1 - S_{n,c}(x),$$

where

$$S_{n,c}(x) := \sum_{k=0}^{\infty}\left(p_{n,k}^{[c]}(x)\right)^2, \quad x \in I_c.$$

The cases $c = -1$, $c = 0$, $c = 1$ correspond, respectively, to the binomial, Poisson, and negative binomial distributions. For other details see also [15], [16].

In this paper we investigate the above entropies with respect to the complete monotonicity.

2 Shannon entropy

A. Let's start with the case $c < 0$.

$H_{n,-1}$ is a concave function; this is a special case of the results of [19]; see also [6], [8], [9] and the references therein.

Here we shall determine the signs of all the derivatives of $H_{n,c}$.

Theorem 1 *Let $c < 0$. Then, for all $k \geq 0$,*

$$H_{n,c}^{(2k+2)}(x) \leq 0, \quad x \in \left(0, -\frac{1}{c}\right), \tag{4}$$

$$H_{n,c}^{(2k+1)}(x) = \begin{cases} \geq 0 & x \in (0, -\frac{1}{2c}], \\ \leq 0 & x \in [-\frac{1}{2c}, -\frac{1}{c}). \end{cases} \tag{5}$$

Proof We have $n = -cl$ with $l \in \mathbb{N}$. As in [10], let us represent $\log(l!)$ by integrals:

$$\log(l!) = \int_0^\infty \left(l - \frac{1-e^{-ls}}{1-e^{-s}}\right) \frac{e^{-s}}{s} ds = \int_0^1 \left(\frac{1-(1-t)^l}{t} - l\right) \frac{dt}{\log(1-t)}. \tag{6}$$

Now using (2), (3) and (6) we get

$$H_{n,c}(x) = H_{l,-1}(-cx) = -l\left[(-cx)\log(-cx) + (1+cx)\log(1+cx)\right] + \int_0^1 \frac{-t}{\log(1-t)} \frac{(1+cxt)^l + (1-t-cxt)^l - 1 - (1-t)^l}{t^2} dt.$$

It is a matter of calculus to prove that

$$\begin{aligned} H''_{n,c}(x) &= cl\left(\frac{1}{x} - \frac{c}{1+cx}\right) \\ &+ c^2 l(l-1) \int_0^1 \frac{-t}{\log(1-t)} \left[(1+cxt)^{l-2} + (1-t-cxt)^{l-2}\right] dt, \end{aligned}$$

and for $k \geq 0$

$$H_{n,c}^{(2k+2)}(x) = cl(2k)! \left(\frac{1}{x^{2k+1}} - \left(\frac{c}{1+cx} \right)^{2k+1} \right)$$
$$+ \quad l(l-1)\dots(l-2k-1)c^{2k+2}$$
$$\int_0^1 \frac{-t}{\log(1-t)} \left[(1+cxt)^{l-2k-2} + (1-t-cxt)^{l-2k-2} \right] t^{2k} dt.$$

For $0 < t < 1$ we have

$$0 < \frac{-t}{\log(1-t)} < 1, \tag{7}$$

so that

$$H_{n,c}^{(2k+2)}(x) \le cl(2k)! \left(\frac{1}{x^{2k+1}} - \left(\frac{c}{1+cx} \right)^{2k+1} \right) + \tag{8}$$

$$+l(l-1)\dots(l-2k-1)c^{2k+2} \int_0^1 \left[(1+cxt)^{l-2k-2} + (1-t-cxt)^{l-2k-2} \right] t^{2k} dt.$$

Repeated integration by parts yields

$$\int_0^1 (1+cxt)^{l-2k-2} t^{2k} dt \le \frac{(2k)!}{(l-2)(l-3)\dots(l-2k-1)(cx)^{2k}} \int_0^1 (1+cxt)^{l-2} dt,$$

and so

$$\int_0^1 (1+cxt)^{l-2k-2} t^{2k} dt \le \frac{(2k)! \left[(1+cx)^{l-1} - 1 \right]}{(l-1)(l-2)\dots(l-2k-1)(cx)^{2k+1}}. \tag{9}$$

Replacing x by $-\frac{1}{c} - x$ we obtain

$$\int_0^1 (1-t-cxt)^{l-2k-2} t^{2k} dt \le \frac{(2k)! \left[1 - (-cx)^{l-1} \right]}{(l-1)(l-2)\dots(l-2k-1)(1+cx)^{2k+1}}. \tag{10}$$

From (8), (9) and (10) it follows that

$$H_{n,c}^{(2k+2)}(x) \le cl(2k)! \left[\frac{(1+cx)^{l-1}}{x^{2k+1}} - \frac{c^{2k+1}(-cx)^{l-1}}{(1+cx)^{2k+1}} \right] \le 0,$$

and this proves (4).

It is easy to verify that $H_{n,c}^{(2k+1)}\left(-\frac{1}{2c}\right) = 0$. Since $H_{n,c}^{(2k+2)} \le 0$, it follows that $H_{n,c}^{(2k+1)}$ is decreasing, and this implies (5).

B. Consider the case $c = 0$.

$H_{n,0}$ is the Shannon entropy of the Poisson distribution. The derivative of this function is completely monotonic: see, e.g., [2, p. 2305]. For the sake of completeness we insert here a short proof.

Theorem 2 *$H'_{n,0}$ is completely monotonic, i.e.,*

$$(-1)^k H_{n,0}^{(k+1)}(x) \geq 0, \quad k \geq 0, \quad x > 0. \tag{11}$$

Proof Let us remark that $H_{n,0}(y) = H_{1,0}(ny)$; so it suffices to investigate the derivatives of $H_{1,0}(x)$.

According to [10, (2.5)],

$$\begin{aligned} H_{1,0}(x) &= x - x\log x + \int_0^\infty \frac{e^{-t}}{t}\left(x - \frac{1-\exp\left(x(e^{-t}-1)\right)}{1-e^{-t}}\right)dt \\ &= x - x\log x - \int_0^1 \left(x - \frac{1-e^{-sx}}{s}\right)\frac{ds}{\log(1-s)}. \end{aligned}$$

It follows that

$$H'_{1,0}(x) = -\log x - \int_0^1 \left(1-e^{-sx}\right)\frac{ds}{\log(1-s)}$$

and for $k \geq 1$,

$$H_{1,0}^{(k+1)}(x) = (-1)^k\left(\frac{(k-1)!}{x^k} + \int_0^1 s^k e^{-sx}\frac{ds}{\log(1-s)}\right). \tag{12}$$

By using (7) we get

$$\int_0^1 \frac{s^k e^{-sx}}{\log(1-s)}ds \geq -\int_0^1 s^{k-1}e^{-sx}ds =$$

$$= -\int_0^x \frac{t^{k-1}}{x^k}e^{-t}dt \geq -\int_0^\infty \frac{1}{x^k}t^{k-1}e^{-t}dt = -\frac{(k-1)!}{x^k}.$$

Combined with (12), this proves (11) for $k \geq 1$. In particular, we see that $H_{n,0}$ is concave and non-negative on $[0, +\infty)$; it follows that $H'_{n,0} \geq 0$ and so (11) is completely proved.

C. Let now $c > 0$.

Theorem 3 *For $c > 0$, $H'_{n,c}$ is completely monotonic.*

Proof Since $H_{m,c}(y) = H_{\frac{m}{c},1}(cy)$, it suffices to study the derivatives of $H_{n,1}(x)$.

By using (2), (3) and

$$\log A = \int_0^\infty \frac{e^{-x} - e^{-Ax}}{x} dx, \quad A > 0,$$

we get

$$H_{n,1}(x) = n\left((1+x)\log(1+x) - x\log x\right) + \int_0^\infty \frac{e^{-ns} - e^{-s}}{s(1-e^{-s})}\left(1 - (1+x-xe^{-s})^{-n}\right) ds$$

$$= n\left((1+x)\log(1+x) - x\log x\right) + \int_0^1 \frac{1-(1-t)^{n-1}}{t\log(1-t)}\left(1-(1+tx)^{-n}\right) dt.$$

It follows that, for $j \geq 1$,

$$\frac{1}{n} H_{n,1}^{(j+1)}(x) = (-1)^{j-1}(j-1)!\left((x+1)^{-j} - x^{-j}\right) +$$

$$+(-1)^{j-1}(n+1)(n+2)\dots(n+j)\int_0^1 \frac{-t}{\log(1-t)}\left[1-(1-t)^{n-1}\right](1+xt)^{-n-j-1}t^{j-1}dt.$$

Using again (7), we get

$$(-1)^{j-1}\frac{1}{n} H_{n,1}^{(j+1)}(x) \leq (j-1)!\left((x+1)^{-j} - x^{-j}\right) +$$
$$+(n+1)(n+2)\dots(n+j)\int_0^1 \left[1-(1-t)^{n-1}\right](1+xt)^{-n-j-1}t^{j-1}dt$$
$$= u(x) + v(x),$$

where

$$u(x) := \frac{(j-1)!}{(x+1)^j} - (n+1)(n+2)\dots(n+j)\int_0^1 t^{j-1}(1-t)^{n-1}(1+xt)^{-n-j-1}dt,$$

$$v(x) := (n+1)(n+2)\dots(n+j)\int_0^1 t^{j-1}(1+xt)^{-n-j-1}dt - \frac{(j-1)!}{x^j}.$$

We shall prove that $u(x) \leq 0$ and $v(x) \leq 0$, $x > 0$. Let us remark that

$$\int_0^1 t^{j-1}(1-t)^{n-1}(1+xt)^{-n-j-1}dt \geq \int_0^1 t^{j-1}(1-t)^n(1+xt)^{-n-j-1}dt, \quad (13)$$

and integration by parts yields

$$\int_0^1 \frac{t^{j-1}(1-t)^n}{(1+xt)^{n+j+1}}dt = \frac{j-1}{(n+1)(x+1)}\int_0^1 \frac{t^{j-2}(1-t)^{n+1}}{(1+xt)^{n+j+1}}dt.$$

Applying repeatedly this formula we obtain

$$\int_0^1 \frac{t^{j-1}(1-t)^n}{(1+xt)^{n+j+1}}dt = \frac{(j-1)!}{(n+1)(n+2)\dots(n+j)}\frac{1}{(x+1)^j}. \quad (14)$$

Now (13) and (14) imply $u(x) \leq 0$.

Using again integration by parts we get

$$\begin{aligned}\int_0^1 t^{j-1}(1+xt)^{-n-j-1}dt &\leq \frac{j-1}{(n+j)x}\int_0^1 t^{j-2}(1+xt)^{-n-j}dt \\ &\leq \dots \leq \frac{(j-1)!}{(n+1)(n+2)\dots(n+j)}\frac{1}{x^j},\end{aligned}$$

which shows that $v(x) \leq 0$.

We conclude that

$$(-1)^{j-1}H_{n,1}^{(j+1)}(x) \leq 0, \quad j \geq 1, x > 0. \quad (15)$$

In particular, (15) shows that $H_{n,1}$ is concave on $[0, +\infty)$; it is also non-negative, which means that $H'_{n,1} \geq 0$. Combined with (15), this shows that $H'_{n,1}$ is completely monotonic, and the proof is finished.

Remark 3.1 (14) *can be obtained alternatively by using the change of variables* $y = (1-t)/(1+xt)$ *and the properties of the Beta function. An alternative proof of the inequality* $v(x) \leq 0$ *follows from*

$$\int_0^1 t^{j-1}(1+xt)^{-n-j-1}dt \leq \frac{1}{x^{j-1}}\int_0^\infty \frac{(xt)^{j-1}}{(1+xt)^{n+j+1}}dt =$$

$$= \frac{1}{x^j}\int_0^\infty \frac{s^{j-1}}{(1+s)^{j+n+1}}ds = \frac{1}{x^j}B(j, n+1) = \frac{1}{x^j}\frac{(j-1)!n!}{(n+j)!}.$$

Corollary 3.1 *The following inequalities are valid for $x > 0$ and $c \geq 0$:*

$$\log \frac{x}{cx+1} \leq \sum_{k=0}^{\infty} p_{n+c,k}^{[c]}(x) \log \frac{k+1}{ck+n} \leq \log \frac{nx+1}{ncx+n}. \tag{16}$$

In particular, for $c = 0$ and $n = 1$,

$$\log x \leq \sum_{k=0}^{\infty} e^{-x} \frac{x^k}{k!} \log (k+1) \leq \log (x+1).$$

Proof We have seen that $H'_{n,c}(x) \geq 0$. An application of (1) yields

$$H'_{n,c}(x) = n \left(\log \frac{1+cx}{x} + \sum_{k=0}^{\infty} p_{n+c,k}^{[c]}(x) \log \frac{k+1}{n+ck} \right).$$

This proves the first inequality in (16); the second is a consequence of Jensen's inequality applied to the concave function $\log t$.

3 Rényi entropy and Tsallis entropy

The following conjecture was formulated in [13]:

Conjecture 3.1 *$S_{n,-1}$ is convex on $[0, 1]$.*

Th. Neuschel [11] proved that $S_{n,-1}$ is decreasing on $\left[0, \frac{1}{2}\right]$ and increasing on $\left[\frac{1}{2}, 1\right]$. The conjecture and the result of Neuschel can be found also in [5].

A proof of the conjecture was given by G. Nikolov [12], who related it with some new inequalities involving Legendre polynomials. Another proof can be found in [4].

Using the important results of Elena Berdysheva [3], the following extension was obtained in [17]:

Theorem 4 *([17, Theorem 9]). For $c < 0$, $S_{n,c}$ is convex on $\left[0, -\frac{1}{c}\right]$.*

A stronger conjecture was formulated in [14] and [17]:

Conjecture 4.1 *For $c \in \mathbb{R}$, $S_{n,c}$ is logarithmically convex, i.e., $\log S_{n,c}$ is convex.*

It was validated for $c \geq 0$ by U. Abel, W. Gawronski and Th. Neuschel [1], who proved a stronger result:

Theorem 5 *([1]). For $c \geq 0$, the function $S_{n,c}$ is completely monotonic, i.e.,*

$$(-1)^m \left(\frac{d}{dx}\right)^m S_{n,c}(x) > 0, \quad x \geq 0, m \geq 0.$$

Consequently, for $c \geq 0$, $S_{n,c}$ is logarithmically convex, and hence convex.

Summing up, for the Rényi entropy $R_{n,c} = -\log S_{n,c}$ and Tsallis entropy $T_{n,c} = 1 - S_{n,c}$, we can state

Corollary 5.1 *i) Let $c \geq 0$. Then $R_{n,c}$ is increasing and concave, while $T'_{n,c}$ is completely monotonic on $[0, +\infty)$.*

ii) $T_{n,c}$ is concave for all $c \in \mathbb{R}$.

Proof

i) Apply Theorem 5.

ii) For $c < 0$, apply Theorem 4. For $c \geq 0$, Theorem 5 shows that $S_{n,c}$ is convex, so that $T_{n,c}$ is concave.

Remark 5.1 *As far as we know, Conjecture 4.1 is still open for $c < 0$, so that the concavity of $R_{n,c}$, $c < 0$, remains to be investigated.*

Acknowledgement

The author is grateful to the referee for valuable comments and very constructive suggestions. In particular, the elegant alternative proofs presented in Remark 1 were kindly suggested by the referee.

References

[1] U. Abel, W. Gawronski, Th. Neuschel, Complete monotonicity and zeros of sums of squared Baskakov functions, Appl. Math. Comput., 258, 130-137 (2015)

[2] J.A. Adell, A. Lekuona and Y. Yu, Sharp bounds on the entropy of the Poisson Law and related quantities, IEEE Trans. Information Theory, 56, 2299-2306 (2010)

[3] E. Berdysheva, Studying Baskakov-Durrmeyer operators and quasi-interpolants via special functions, J. Approx. Theory, 149, 131-150 (2007)

[4] I. Gavrea, M. Ivan, On a conjecture concerning the sum of the squared Bernstein polynomials, Appl. Math. Comput., 241, 70-74 (2014)

[5] H. Gonska, I. Raşa, M.-D. Rusu, Chebyshev-Grüss-type inequalities via discrete oscillations, Bul. Acad. Stiinte Repub. Mold. Mat., 1, (74), 63-89; arxiv 1401.7908 [math.CA] (2014)

[6] P. Harremoës, Binomial and Poisson distributions as maximum entropy distributions, IEEE Trans. Information Theory, 47, 2039 - 2041 (2001)

[7] M. Heilmann, Erhöhung der Konvergenzgeschwindigkeit bei der Approximation von Funktionen mit Hilfe von Linearkombinationen spezieller positiver linearer Operatoren, Habilitationsschrift, Universität Dortmund (1992)

[8] E. Hillion, Concavity of entropy along binomial convolutions, Electron. Commun. Probab., 17, 1-9 (2012)

[9] E. Hillion, O. Johnson, A proof of the Shepp-Olkin entropy concavity conjecture, arXiv: 1503.01570v1, (2015)

[10] C. Knessl, Integral representations and asymptotic expansions for Shannon and Rényi entropies, Appl. Math. Lett., 11, (1998), 69-74.

[11] Th. Neuschel, Unpublished manuscript (2012)

[12] G. Nikolov, Inequalities for ultraspherical polynomials. Proof of a conjecture of I. Raşa, J. Math. Anal. Appl., 418, (2014), 852-860.

[13] I. Raşa, Unpublished manuscripts (2012)

[14] I. Raşa, Special functions associated with positive linear operators, arxiv: 1409.1015v2, (2014)

[15] I. Raşa, Rényi entropy and Tsallis entropy associated with positive linear operators, arxiv: 1412.4971v1, (2014)

[16] I. Raşa, Entropies and the derivatives of some Heun functions, arxiv: 1502.05570v1, (2015)

[17] I. Raşa, Entropies and Heun functions associated with positive linear operators, Appl. Math. Comput., 268, 422-431 (2015)

[18] A. Rényi, On measures of entropy and information, Proc. Fourth Berkeley Symp. Math. Statist. Prob., Vol. 1, Univ. of California Press, pp. 547-561 (1961)

[19] L.A. Shepp, I. Olkin, Entropy of the sum of independent Bernoulli random variables and of the multinomial distribution, Proc. Contributions to Probability, New York, pp. 201-206 (1981)

[20] C. Tsallis, Possible generalization of Boltzmann-Gibbs statistics, J. Stat. Phys., 52, 479-487 (1988)

[21] M. Wagner, Quasi-Interpolaten zu genuinen Baskakov-Durrmeyer-Typ Operatoren, Shaker Verlag, Aachen (2013)